

BIBLIOGRAFIA

- Aleotti, P., Chowdhury, R., 1999. Landslide hazard assessment: summary review and new perspectives. Bull Eng Geol Env. 58, 21-44.
- Antonioli, F., Kershaw, S., Rust, D., Verrubbi, V., 2003. Holocene sea-level change in Sicily and its implications for tectonic models: new data from the Taormina area, northeast Sicily. Marine Geol. 3293, 1-19.
- APAT, Servizio Geologico d'Italia, Dipartimento Difesa del Suolo: Carta Geologica d'Italia alla scala 1:50.000, Foglio 601, Messina. Reggio di Calabria, S.EL.CA, Firenze, 2008.
- AA. VV., 2000. Carta Geologica della provincia di Messina, Foglio 1, Scala 1:50.000. S.EL.CA., Fiernze.
- Agliardi, F., Zanchi, A., Crosta, G.B., 2009. Tectonic vs. gravitational morphostructures in the central Eastern Alps (Italy): Constraints on the recent evolution of the mountain range, Tectonophys. 474, 250-270.
- Agnesi, V., Rasà, R., Puglisi, C., Gioè, C., Privitera, B., Cappadonia, C., Conoscenti, C., Pino, P., Rotigliano, E., 2009. La franosità diffusa dell'1 Ottobre 2009 nel territorio ionico-peloritano della Provincia di Messina: stato delle indagini e prime considerazioni sulle dinamiche geomorfiche attivate. Geologi di Sicilia 4, 23-30.
- Aleotti, P., Chowdhury, R., 1999. Landslide hazard assessment: summary review and new perspectives. Bull. Eng. Geol. Env. 58, 21-44.
- Amodio-Morelli, L., Bonardi, G., Colonna, V., Dietrich, D., Giunta, G., Ippolito, F., Liguori, V., Lorenzoni, S., Paglionico, A., Perrone, V., Piccarreta, G., Russo, M., Scandone, P., Zanettin-Lorenzoni, E., Zuppetta, A., 1976. L'Arco Calabro-peloritano nell'orogene appenninico-maghrebide, Mem. Soc. Geol. It. 17, 1-60.
- Antonioli, F., Ferranti, L., Fontana, A., Amorosi, A., Bondesan, A., Braitenberg, C., Dutton, A., Fontolan, G., Furlani, S., Lambeck, K., Mastronuzzi, G., Monaco, C., Spada, G., Stocchi, P.,

2009. Holocene relative sea-level changes and vertical movements along the Italian and Istrian coastlines, *Q. Int.* 206, 102-133.
- Antonioli, F., Ferranti, L., Lambeck, K., Kershaw, S., Verrubbi, V., Pra, G.D., 2006. Late Pliestocene to Holocene record of changing uplift rates in southern Calabria and northeastern Sicily (southern Italy, Central Mediterranean Sea). *Tectonophys.* 422, 23-40.
- Antonioli, F., Kershaw, S., Rust, D., Verrubbi, V., 2003. Holocene sea-level change in Sicily and its implications for tectonic models: new data from the Taormina area, northeast Sicily, *Mar. Geol.* 196, 53-71.
- APAT, Servizio Geologico d'Italia, Dipartimento Difesa del Suolo, 2008. Note illustrative della Carta Geologica d'Italia alla scala 1:50.000, Foglio 601 Messina. Reggio di Calabria, S.E.L.CA, Firenze, 2008.
- APAT, Servizio Geologico d'Italia, Dipartimento Difesa del Suolo, 2010. Note illustrative della Carta Geologica d'Italia alla scala 1:50.000, Foglio 613 Taormina, S.E.L.CA, Firenze, 2010
- Argnani, A., Brancolini, G.; Bonazzi, C., Rovere, M., Accaino, F.; Zgur, F., Lodolo, E., 2006. The results of the Taormina 2006 seismic survey: Possible implications for active tectonics in the Messina Straits. *Tectonophys.* 476, 159-169.
- Aufmuth, R.E., 1973. A systematic determination of engineering criteria for rocks. *Bull. Assoc. Eng. Geol.* 11, 235–245.
- Avena, G.C., Giuliano, G., Lupia Palmieri, E., 1967. Sulla valutazione quantitativa della gerarchizzazione ed evoluzione dei reticolii fluviali., *Boll. Soc. Geol. It.* 86, 781-796.
- Aydin, A., 2009. ISRM Suggested method for determination of the Schmidt hammer rebound hardness: Revised version. *Int. J. Rock Mech. Min. Sci.* 46, 627–634.
- Aydin, A., Basu, A., 2005. The Schmidt hammer in rock material characterization *Eng. Geol.* 81, 1-14.

- Baeza, C., Corominas, J., 2001. Assessment of shallow landslide susceptibility by means of multivariate statistical techniques. *Earth Process. Surf. Landforms* 26/12, 1251–1263.
- Bai, T., Maerten, L., Gross, M.R., Aydin, A., 2002. Orthogonal cross joints: do they imply a regional stress rotation?. *J. Struct. Geol.* 24, 77-88.
- Basile, G., 2010. Evento del 1 Ottobre 2009, Carta dei dissesti, Dipartimento Regionale della Protezione Civile, Sicilia.
- Basu A., Aydin, A., 2004. A method for normalization of Schmidt hammer rebound values. *Int. J. Rock Mech. Min. Sci.* 41, 1211-1214.
- Bell, J.S., 1996. In situ stresses in sedimentary rocks (part 2): applications of stress measurements, *Geosci. Canada* 23(3), 135–153
- Beverly, B.E., Schoenwolf, D.A., Brierly, G.S., 1979. Correlations of rock index values with engineering properties and the classification of intact rock.
- Bianca, M., Catalano, S., De Guidi, G., Gueli, A.M., Monaco, C., Ristuccia, G.M., Stella, G., Tortorici, G., Tortorici, L., Troja, S.O., 2011. Luminescence chronology of Pleistocene marine terraces of Capo Vaticano peninsula (Calabria, Southern Italy). *Q. Int.* 232, 114-121.
- Bianca, M., Monaco, C., Tortorici, L., Cernobori, L., 1999. Quaternary normal faulting in southeastern Sicily (Italy): a seismic source for the 1693 large earthquake. *Geophys. J. Int.* 130, 370-394.
- Bieniawski, Z., T., 1979. The geomechanics classification in rock engineering applications. Proceedings of 4th International Congress on Rock Mechanics, 2-8 September 1979. Montreux, Switzerland.
- Billi, A., Presti, D., Faccenna, C., Neri, G., Orecchio, B., 2007. Seismotectonics of the Nubia plate compressive margin in the south Tyrrhenian region, Italy: Clues for subduction inception. *J. Geophys. Res.* 112, B08302.

Bibliografia

- Blahut, J., van Westen, C.J., Sterlacchini, S., 2010. Analysis of landslide inventories for accurate prediction of debris-flow source areas, *Geomorphology* 119, 36-51.
- Boccaletti, M., Nicolich, R., Tortorici, L., 1984. The Calabrian Arc and the Ionian Sea in the dynamic evolution of the central Mediterranean. *Marin. Geol.* 55, 219-245.
- Boccaletti, M., Nicolich, R., Tortorici, L., 1990. New data and hypothesis on the development of the Tyrrhenian basin. *Palaeogeography, Palaeoclimatology, Palaeoecology* 77, 15-40.
- Bonardi, G., de Capoa, P., Di Staso, A., Martin-Martin, M., Martin-Rojas, I., Perrone, V., Tent-Manclus, J.E., 1982. New constraints to the geodynamic evolution of the southern sector of the Calabria-Peloritani Arc (Italy), *C. R. Geosci.* 334, 423-430.
- Bonfiglio, L., Violanti, D., 1983. Prima segnalazione di Tirreniano ed evoluzione pleistocenica del Capo Peloro (Sicilia nord-orientale). *Geogr. Fis. Dinam. Quat.* 6, 3-15.
- Borrelli, L., Greco, R., Gullà, G., 2007. Weathering grade of rock masses as a predisposing factor to slope instabilities: Reconnaissance and control procedures, *Geomorphology* 84, 158-175.
- Brenning, A., 2005. Spatial prediction models for landslide hazards: review, comparison and evaluation, *Nat. Haz. Earth Syst. Sci.* 5, 853-862.
- Burbank, D.W., 2002. Rates of erosion and their implications for exhumation, *Mineralogical Magazine*, 66(1), 25-52.
- Burbank, D.W., Pinter, N., 1999. Landscape evolution: the interactions of tectonics and surface processes, *Basin Res.*, 11, 1-6.
- Calcaterra, D., Parise, M., 2005. Landslide types and their relationships with weathering in a Calabrian basin, southern Italy, *Bull. Eng. Geol. Environ.* 64, 193-207.
- Cargill, J.S., Shakoor, A., 1990. Evaluation of empirical methods for measuring the uniaxial compressive strength. *Int. J. Rock. Mech. Min. Sci.* 27, 495-503.

Bibliografia

- Caporali; A., Aichhorn, C., Barlik, M., Becker, M., Fejes, I., Gerhatova, L., Ghitau, D., Grenerczy, G., Hefty, J., Krauss, S., Medak, D., Milev, G., Mojzes, M., Mulic, M., Nardo, A., Pesec, P., Rus, T., Simek, J., Sledzinski, J., Solaric, M., Stangl, G., Stopar, M., Vespe, F., Virag, G., 2009. Surface kinematics in the Alpine–Carpathian–Dinaric and Balkan region inferred from a new multi-network GPS combination solution. *Tectonophys.* 474, 295–321.
- Caputo, R., 1991. Comparison between joints and faults as brittle structures used for evaluating stress field. *Annales Tectonicae* 5(1), 74-84.
- Caputo, R., 2005. Stress variability and brittle tectonic structures. *Earth. Sci. Rev.* 70, 103-127.
- Caputo, R., 2010. Why joints are more abundant than faults. A conceptual model to estimate their ratio in layered carbonate rocks. *J. Struct. Geol.* 32, 1257-1270.
- Caputo, M., Caputo, R., 1989. Estimate of the regional stress field using joint system. *Bull. Geol. Soc. Greece*, 23(1), 101-118
- Carminati, E., Wortel, M.J.R., Spakman, W., Sabadini, R., 1998. The role of slab detachment processes in the opening of the western–central Mediterranean basins: some geological and geophysical evidence. *Earth Planet. Sci. Lett.* 160, 651-665.
- Carrara, A., Cardinali, M., Detti, R.; Guzzetti, F., Pasqui, V., Reichenbach, P., 1991. GIS techniques and statistical models in evaluating landslide hazard, *Earth Process. Surf. Landforms* 16, 427-445.
- Carrara, A., Guzetti, F., Cardinali, M., Reichenbach, P., 1999. Use of GIS Technology in the Prediction and Monitoring of Landslide Hazard, *Nat. Hazards* 20, 117-135.
- Cascini, L., 2008. Applicability of landslide susceptibility and hazard zoning at different scales, *Eng. Geol.* 102, 164-177.
- Catalano, S., Cinque, A., 1995. Dati preliminari sull'evoluzione neotettonica dei Peloritani settentrionali (Sicilia nordorientale) sulla base dei dati morfologici. *Studi Geol. Camerti*, Vol. Spec. 2, 113–123.

Bibliografia

- Catalano, S., De Guidi, G., 2003. Late Quaternary uplift of northeastern Sicily: relation with the active normal faulting deformation. *J. Geodyn.* 36, 445-467.
- Catalano, S., De Guidi, G., Monaco, C., Tortorici, G., Tortorici, L., 2003. Long-term behaviour of the late Quaternary normal faults in the Straits of Messina area (Calabrian arc): structural and morphological constraints. *Q. Int.* 101/102, 81-91.
- Catalano, S., De Guidi, G., Monaco, C., Tortorici, G., Tortorici, L., 2008. Active faulting and seismicity along the Siculo-Calabrian Rift Zone (southern Italy). *Tectonophys.* 453, 177-192.
- Catalano, S., Di Stefano, A., 1997. Sollevamento e tettogenesi Pleistocenica lungo il margine tirrenico dei Monti Peloritani: integrazione dei dati geomorfologici, strutturali e biostratigrafici. *Il Quaternario* 10, 337-342.
- Cavazza, W., Forese, C.W., 2003. The Mediterranean region—a geological primer. *Episodes*, 26(3), 160-168
- Cevik, E., Topal, T., 2003. GIS-based landslide susceptibility mapping for a problematic segment of the natural gas pipeline, Hendek (Turkey). *Environ. Geol.* 44, 949-962.
- Chung, C.F., Fabbri, A.G., 2003. Validation of Spatial Prediction Models for Landslide Hazard Mapping, *Nat. Hazards* 30, 451-472.
- Cifelli, F., Rossetti, F., Mattei, M., Hirt, A.M., Funiciello, R., Tortorici, L., 2004. An AMS, structural and paleomagnetic study of quaternary deformation in eastern Sicily. *J. Struct. Geol.* 26, 29-46.
- Cirrincione, R., Fazio, E., Ortolano, G., Pezzino, A., Punturo, R., 2012. Fault-related rocks: deciphering the structural–metamorphic evolution of an accretionary wedge in a collisional belt, NE Sicily. *Int. Geol. Rev.* 54(8), 94-95.
- Coe, J.A., Godt, J.W., Wait, T.C., and Kean, J.W., 2007, Field reconnaissance of debris flows triggered by a July 21, 2007, thunderstorm in Alpine, Colorado, and vicinity: U.S. Geological Survey Open File Report 2007-1237, 25 p.

Bibliografia

- Coelho-Netto, A.L., Avelar, A.S., Fernandes, M.C., Lacerda, W.A., 2007. Landslide susceptibility in a mountainous geoecosystem, Tijuca Massif, Rio de Janeiro: The role of morphometric subdivision of the terrain. *Geomorphology* 87, 120-131.
- Cruden, D.M., and Varnes, D.J., 1996. Landslide types and processes. In: Turner, A.K., and Schuster, R.L. (eds), *Landslides investigation and mitigation, special report 247*. Transportation Research Board, National Academy Press, Washington D.C: 36-75.
- Cyr, A.J., Granger, D.E., Olivetti, V., Molin, P., 2010. Quantifying rock uplift rates using channel steepness and cosmogenic nuclide-determined erosion rates: Examples from northern and southern Italy. *Lithosphere* 2/3, 188-198.
- D'Agostino N, Avallone A, Cheloni D, D'Anastasio E, Mantenuto S, Selvaggi G 2008 Active tectonics of the Adriatic region from GPS and earthquake slip vectors. *J Geophy Res* 113, B12413
- D'Agostino, N., Selvaggi, G., 2004. Crustal motion along the Eurasia-Nubia boundary in the Calabrian Arc and Sicily and active extension in the Messina Straits from GPS measurements. *J. Geophy. Res.* 109, B11402
- Dahal, R.K., Hasegawa, S., Nonomura, A., Yamanaka, M., Dhakal, S., Paudyal, P., 2008. Predictive modelling of rainfall-induced landslide hazard in the Lesser Himalaya of Nepal based on weights-of-evidence. *Geomorphology* 102, 496-510.
- Dahal, R.K., Hasegawa, S., Nonomura, A., Yamanaka, M., T., Masuda, Nishino, K., 2008. GIS-based weights-of-evidence modelling of rainfall-induced landslides in small catchments for landslide susceptibility mapping. *Environ. Geol.* 54, 311-324.
- Deb, S.K., El-Kady, A.I., 2009. Susceptibility assessment of shallow landslides on Oahu, Hawaii, under extreme-rainfall events, *Geomorphology* 108, 129-233.
- De Gregorio, S., Rotolo, S.G., Villa, I.M., 2003. Geochronology of the medium to high-grade metamorphic units of the Peloritani Mts., Sicily. *Int. J. Earth Sci. (Geol. Rundsch.)* 92, 852-872.

Bibliografia

- De Guidi, G., Catalano, S., Monaco, C., Tortorici, L., 2003. Morphological evidence of Holocene coseismic deformation in the Taormina region (NE Sicily). *J. Geodyn.* 36, 193-211.
- De Guidi, G., Scudero, S., Landslide susceptibility assessment in the Peloritani Mts. (Sicily, Italy) and clues for tectonic control of relief processes. *Nat. Hazards Earth Syst. Sci.* *In press*
- Del Ventisette, C., Garfagnoli, F., Ciampalini, A., Battistini, A., Gigli, G., Moretti, S., Casagli, N., 2012. An integrated approach to the study of catastrophic debris-flows: geological hazard and human influence. *Nat. Hazards Earth Syst. Sci.* 12, 2907-2922.
- De Vita, P., Reichenbach, P., 1998. Rainfall-triggered landslides: a reference list. *Environ. Geol.* 35 (2–3), 219-233.
- Densmore, A.L., Ellis, M.A., Anderson, R.S., 1998. Landsliding and the evolution of normal-fault-bounded mountains. *J. Geophy. Res.*, 103 (B7), 15203-15219.
- Dewey, J.F., Helman, M.L., Knott, S.D., Turco, E., Hutton, D.H.W., 1989. Kinematics of the western Mediterranean. *Geol. Soc. London, Special Publications* 45, 265-283.
- Dietrich, W.E., Dunne, T., Humphrey, N.F., Reid, L.M., 1982. Construction of sediment budgets for drainage basins. In: *Sediment Budgets and routing in forested drainage basins, Proceedings of the Symposium; 31 May - 1 June 1982; Corvallis, Oregon. Gen. Tech. Rep. PNW-141*. Portland, Oregon: Pacific Northwest Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture, 5-23.
- Di Stefano, A., Longhitano, S.G., 2009. Tectonics and sedimentation of the Lower and Middle Pleistocene mixed siliciclastic/bioclastic sedimentary successions of the Ionian Peloritani Mts (NE Sicily, Southern Italy): the onset of opening of the Messina Strait. *Cent. Eur. J. Geosci.* 1(1), 33-62.
- Dunne, W.M., Hancock, P.L., 1994. Palaeostress analysis of small-scale brittle structures. In: Hancock, P. (Editor), *Continental deformation*, Pergamon Press.
- Engelder, T., 1987. Joints and shear fractures in rock. In: Atkinson, B. (Editor), *Fracture Mechanics of Rock*. Academic Press, Orlando, 27-69.

Bibliografia

- Faccenna, C., Molin, P., Orecchio, B., Olivetti, B., Bellier, O., Funicello, F., Minelli, L., Piromallo, C., Billi, A., 2011. Topography of the Calabria subduction zone (southern Italy): Clues for the origin of Mt. Etna. *Tectonics* 30, TC1003.
- Faccenna, Thorsten W. Becker, Francesco Pio Lucente,3 Laurent Jolivet and Federico Rossetti, 2001a. History of subduction and back-arc extension in the Central Mediterranean. *Geophys. J. Int.* 145, 809-820.
- Faccenna; C., Funicello, F., Giardini, D., Luente, P., 2001b. Episodic back-arc extension during restricted mantle convection in the Central Mediterranean. *Earth Planet. Sci. Lett.* 187, 105-116.
- Falaschi, F., Giacomelli, F., Federici, P.R., Puccinelli, A., D'Amato Avanzi, G., Pochini, A., Ribolini, A., 2009. Logistic regression versus artificial neural networks: landslide susceptibility evaluation in a sample area of the Serchio River valley, Italy, *Nat. Hazards* 50, 551-569.
- Falconi, L., Campolo, D., Leoni, G., Lumaca, S., Puglisi, C.: Geomorphology hazard assessment of Giampilieri and Briga river basins after the rainfall event on the October 1, 2009 (Sicily, Italy), The Second World Landslide Forum, “Putting science into practice”. Global Promotion Committee of the International Programme on Landslides (IPL), Rome, Italy, 2011.
- Federici, P.R., Puccinelli, A., Cantarelli, E., Casarosa, N., D'Amato Avanzi, G., Falaschi, F., Giannecchini, R., Pochini, A., Ribolini, A., Bottai, M., Salvati, N., Testi, C., 2007. Multidisciplinary investigations in evaluating landslide susceptibility. An example in the Serchio River valley (Italy), *Q. Int.* 171-172, 52-63.
- Fell, R., Corominas, J., Bonnard, C., Cascini, L., Leroi, E., Savage, W.Z., on behalf of the JTC-1 Joint Technical Committee on Landslides and Engineered Slopes, 2008. Guidelines for landslide susceptibility, hazard and risk zoning for land use planning, *Eng. Geol.* 102, 85-98.
- Festa, V., Messina, A., Paglionico, A., Piccarreta, G., Rottura, A., 2004. Pre-Triassic history recorded in the Calabria–Peloritani segment of the Alpine chain, southern Italy. An overview. Spec Iss 2: a showcase of the Italian research in metamorphic petrology. *Period. Mineral.* 73, 57-71.

- Firth, C., Stewart, I., Mc Guire, W.J., Kershaw, S., Vita-Finzi, C., 1996. Coastal elevation changes in eastern Sicily: implications for volcano in stability at Mount Etna. Volcano instability on the earth and other planets. Geol. Soc. London Spec. Pub. 110, 153-167.
- Font, M., Amorese, D., Lagarde, J.L., 2010. DEM and GIS analysis of the stream gradient index to evaluate effects of tectonics: The Normandy intraplate area (NW France). Geomorphology 119, 172-180.
- García-Rodríguez, M.J., Malpica, J.A., Benito, B., Díaz, M., 2008. Susceptibility assessment of earthquake-triggered landslides in El Salvador using logistic regression. Geomorphology 95, 172-191.
- Glade, T., 1998. Establishing the frequency and magnitude of landslide-triggering rainstorm events in New Zealand, Environ. Geol. 35 (2/3), 160-174.
- Glade, T., Crozier, M., Smith, P., 2000. Applying Probability Determination to Refine Landslide-triggering Rainfall Thresholds Using an Empirical “Antecedent Daily Rainfall Model”. Pure Appl. Geophys. 157, 1059-1079.
- Ghisetti, F., 1979. Relazioni tra strutture e fasi trascorrenti e distensive lungo i sistemi Messina-Fiumefreddo, Tindari-Letojanni e Alia-Malvagna (Sicilia Nord-orientale): uno studio microtetonico. Geologica Romana 18, 23-58.
- Ghisetti, F., 1992. Fault parameters in the Messina Strait (southern Italy) and relations with the seismogenic source. Tectonophys. 201, 117-133.
- Giunta, G., Gueli, A.M., Monaco, C., Orioli, S.; Ristuccia, G.M., Stella, G., Troja, S.O., 2012. Middle-Late Pleistocene marine terraces and fault activity in the Sant'Agata di Militello coastal area (north-eastern Sicily). J. Geodyn. 55, 32-40.
- Giunta, G., Nigro, F., 1999. Tectono-sedimentary constraints to the Oligocene-to-Miocene evolution of the Peloritani thrust belt (NE Sicily). Tectonophys. 315, 287-299.

Bibliografia

- Godt, J.W., Baum R.L., Savage, W.Z., Salciarini, D., Schultz, W.H., Harp, E.L., 2008. Transient deterministic shallow landslide modeling: Requirements for susceptibility and hazard assessments in a GIS framework, *Eng. Geol.* 102, 214-236.
- Goes S, Giardini D, Jenny S, Hollenstein C, Kahle HG, Geiger,A. 2004 A recent reorganization in the south-central Mediterranean. *Earth Planet Sci. Lett.* 226, 335-345.
- Goldsworthy, M., Jackson, J., 2000. Active normal fault evolution in Greece revealed by geomorphology and drainage patterns, *J. Geol. Soc. London*, 157, 967-981.
- Gomez, H., Kavzoglu, T., 2005. Assessment of shallow landslide susceptibility using artificial neural networks in Jabonosa River Basin, Venezuela. *Eng. Geol.* 78, 11-27.
- Goswami, R., Brocklehurst, S.H, Mitchell, N.C., 2012. Erosion of a tectonically uplifting coastal landscape, NE Sicily, Italy, *Geomorphology*, 171-172, 114-126.
- Goswami, R., Mitchell, N.C., Brocklehurst, S.H., 2011. Distribution and causes of landslides in eastern Peloritani of NE Sicily and western Aspromonte of SW Calabria, Italy, *Geomorphology*, 132, 111-122.
- Grande, A., Di Vincenzo, G., Prosser, G., Caggianelli, A., 2009. Direct evidence of Middle Oligocene extension in the Calabria–Peloritani terrane from co-seismic faulting: the pseudotachylite-bearing shear zones of Palmi (southern Calabria, Italy). *Terra Nova* 21, 293-303
- Greco, R., Sorriso-Valvo, M., 2005. Relationships between joint apparent separation, Schmidt hammer rebound value, and distance to faults, in rocky outcrops, Calabria, Southern Italy, *Eng. Geol.* 78, 309-320.
- Guarnieri, P., Pirrotta, C., 2008. The response of drainage basins to the late Quaternary tectonics in the Sicilian side of the Messina Strait (NE Sicily), *Geomorphology* 95, 260-273.
- Gueguen, E., Doglioni, C., Fernandez, M., 1998. On the post-25 Ma geodynamic evolution of the western Mediterranean. *Tectonophysics* 298, 259-269

Bibliografia

- Gullà, G., Antronico, L., Iaquinta, P., Terranova, O., 2008. Susceptibility and triggering scenarios at a regional scale for shallow landslides, *Geomorphology*, 99, 39-58.
- Gunther, A., Thiel, C., 2009. Combined rock slope stability and shallow landslide susceptibility assessment of the Jasmund cliff area (Rugen Island, Germany), *Nat. Hazards Earth Syst. Sci.* 9, 687-698.
- Guzzetti, F., Carrara, A., Cardinali, M., Reichenbach, P., 1999. Landslide hazard evaluation: a review of current techniques and their application in a multi-scale study, Central Italy, *Geomorphol.* 31, 181-216.
- Guzzetti, F., Galli, M., Reichenbach, P., Ardizzone, F., Cardinali, M., 2006a. Landslide hazard assessment in the Collazzone area, Umbria, Central Italy. *Nat. Haz. Earth System Sci.* 6, 115-131.
- Guzzetti, F., Reichenbach, P., Ardizzone, F., Cardinali, M., Galli, M., 2006b. Estimating the quality of landslide susceptibility models, *Geomorphology*, 81, 166-184.
- Guzzetti, F., Stark, C.,P., Salvati, P., 2005. Evaluation of Flood and Landslide Risk to the Population of Italy. *Environmental Management* 36(1), 15-36.
- Hack, J.T., 1973. Stream-profile analysis and stream-gradient index. *U. S. Geol. Surv. J. Res.* 1, 421-429.
- Hancock, P.L., Al-Kadhi, A., Barka, A.A., Bevan, T.G, 1987. Aspect of analyzing brittle structures. *Annales Tectonicae* 1, 5-19
- Haramy K.Y., DeMarco M.J., 1985. Use of Schmidt hammer for rock and coal testing. *Proceedings of 26th US symposium on rock mechanics* 26–28 June, Rapid City. Balkema, Rotterdam, 549-555.
- Harp, H.L., Reid, M.E., McKenna, J.P., Michael, J.A., 2009. Mapping of hazard from rainfall-triggered landslides in developing countries: Examples from Honduras and Micronesia, *Eng. Geol.* 104, 295-311.

Bibliografia

- Heimsath, A.M., Dietrich, W.E., Nishiizumi, K., Finkel, R.C., 1999. Cosmogenic nuclides, topography, and the spatial variation of soil depth. *Geomorphology* 27, 151-172.
- Hengl, T., 2006. Finding the right pixel size, *Computers & Geosciences* 32, 1283-1298.
- Heymes, T., Monié, P., Arnaud, N., Pêcher, A., Bouillin, JP., Compagnoni, R., 2010. Alpine tectonics in the Calabrian–Peloritan belt (southern Italy): New $^{40}\text{Ar}/^{39}\text{Ar}$ data in the Aspromonte Massif area, *Lithos* 114, 461-472.
- Highland, L.M., and Bobrowsky, Peter, 2008, The landslide handbook—A guide to understanding landslides: Reston, Virginia, U.S. Geological Survey Circular 1325, 129 p.
- Hren, M.T, Hilley, G.E., Chamberlain, C.P., 2007. The relationship between tectonic uplift and chemical weathering rates in the Washington Cascades: field measurements and model predictions, *Am. J. Sci.*, 307, 1041–1063.
- Hungr, O., Evans, S.G, Bovis, M.J., Hutchinson, J.N., 2001. A review of the classification of landslides of the flow type. *Environmental & Engineering Geoscience* 7(3), 221-238.
- Hungr, O., McDougall, S., Wise, M., Cullen, M., 2008. Magnitude–frequency relationships of debris flows and debris avalanches in relation to slope relief. *Geomorphology* 96, 355–365.
- Hutchinson, J.N., 1988. General report: morphological and geotechnical parameters of landslides in relation to geology and hydrology. 5th International Symposium on Landslides, Balkema, Rotterdam, 1, 3-35.
- Ibetsberger, H.J., 1996. The Tsergo Ri landslide: an uncommon area of high morphological activity in the Langthang valley, Nepal. *Tectonophys.* 260, 85-93.
- Ietto, F., Donato, F.F., Ietto, A., 2007. Recent reverse faults and landslides in granitoid weathered profiles, Serre Mountains (southern Calabria, Italy). *Geomorphology* 87, 196-206.
- ISRM, International Society on Rock Mechanics, 1981. "Basic geotechnical description of rock masses." *International Journal of Rock Mechanics, Mining Sciences, and Geomechanics Abstracts*, 18,85-110.

Bibliografia

- Irfan, T.Y., Dearman, W.R., 1978. Engineering classification and index properties of a weathered granite. Bull Assoc Eng. Geol. 17, 79-90.
- Kahraman, S., 2001. Evaluation of simple methods for assessing the uniaxial compressive strength of rock. Int. J. Rock Mech. Min. Sci. 38, 981-994.
- Keefer, D.K., 1984. Landslides caused by earthquakes. Bull. Geol. Soc. Am. 95, 406-421.
- Keefer, D.K., 2000. Statistical analysis of an earthquake-induced landslide distribution —the 1989 Loma Prieta, California event. Eng. Geol. 58, 231-249.
- Keefer, D.K., 2002. Investigating landslides caused by earthquakes—a historical review. Surveys in Geophysics 23, 473-510.
- Keller, E.A., Pinter, N., 1996. Active Tectonics: Earthquakes, Uplift, and Landscape, Prentice Hall, Upper Saddle River, New Jersey, pp. 338.
- Kidybinski, 1980. Bursting liability indices of coal. Int. J. Rock Mech. Min. Sci. Geomech. Abstr. 17, 167-171.
- King, G.C.P., Stein, S.S., Rundle, J.B., 1988. The growth of geological structures by repeated earthquakes 1. Conceptual framework. J. Geophys. Res. 93 (B11), 13,307-13,318.
- Korup, O., Clague, J.J., Hermanns, R.L., Hewitt, K., Strom, A.L., Weidinger, J.T., 2007. Giant landslides, topography, and erosion, Earth Pl. Sci. Lett., 261, 578-589.
- Korup, O., Densmore, A.L. Schlunegger, G., 2010. The role of landslides in mountain range evolution, Geomorphology, 102, 77-90.
- Lambeck, K., Antonioli, F., Purcell, A., Silenzi, S., 2004. Sea-level change along the Italian coast for the past 10,000 yr, Quaternary Sci. Rev. 23, 1567-1598.
- Lee, S., Min, K., 2001. Statistical analysis of landslide susceptibility at Yongin, Korea. Environ. Geol. 40, 1095-1113.

Bibliografia

- Le Pera, E., Sorriso-Valvo, M., 2000. Weathering and morphogenesis in a mediterranean climate, Calabria, Italy, *Geomorphology*, 34, 251-270.
- Leroueil, S., Vaunat, J., Picarelli, L., Locat, J., Lee, H., and Faure, R.: 1996, Geotechnical characterisation of slope movements, Proceedings 7th International Symposium on Landslides, Trondheim. A. A. Balkema, Rotterdam. Vol. 1, 53-74.
- Malinverno, A., Tyan, W.B.F., 1986. Extension in the Tyrrhenian Sea and shortening in the Apennines as result of arc migration driven by sinking of the lithosphere. *Tectonics* 5/2, 227-245.
- Mandl, G., 2005. Rock Joints. The mechanical genesis. Springer-Verlag Berlin Heidelberg
- Marques, E.A.G., Barroso, E.V., Menez Filho, A.P., Vargas Jr., E.A., 2010. Weathering zones on metamorphic rocks from Rio de Janeiro—Physical, mineralogical and geomechanical characterization, *Eng. Geol.* 111, 1-18.
- Mauffret, A., 2007. The Northwestern (Maghreb) boundary of the Nubia (Africa) Plate. *Tectonophys.* 429, 21-44.
- Maugeri, M., Motta, E., 2010. Effects of Heavy Rainfalls on Slope Behaviour: The October 1, 2009 Disaster of Messina (Italy). *Geotechnics/Earthquake Geotechnics towards Global Sustainability*, 12-14 January, 2010. Kyoto, Japan.
- McClusky, S., Reilinger, R., Mahmoud, S., Ben Sari, D., Tealeb, A., 2003. GPS constraints on Africa (Nubia) and Arabia plate motions. *Geophys. J. Int.* 155, 126-138.
- Miles, S.B., Keefer, D.K., 2009. Evaluation of CAMEL — comprehensive areal model of earthquake-induced landslides, *Engineering Geology* 104, 1-15.
- Miller, S., Brewer, T., Harris, N., 2009. Rainfall thresholding and susceptibility assessment of rainfall-induced landslides: application to landslide management in St Thomas, Jamaica. *Bull. Eng. Geol. Environ.* 68, 539-550.

Bibliografia

- Monaco, C., Antonioli, F., De Guidi, G., Lambeck, K., Tortorici, L., Verrubbi, V., 2004. Holocene tectonic uplift of the Catania Plain (Eastern Sicily). *Quat. Nov.* 8, 171-185.
- Monaco, C., Bianca, M., Catalano, S., De Guidi, G., Tortorici, L., 2002. Sudden change in the Late Quaternary tectonic regime in eastern Sicily: evidences from geological and geomorphological features. *Boll. Soc. Geol. It. Vol. Spec.* 1, 901-913.
- Monaco, C., Tapponier, P., Tortorici, L., Gyllot, P.Y., 1997. Late Quaternary slip rates on the Acireale-Piedimonte normal faults and tectonic origin of Mt. Etna (Sicily). *Earth Planet. Sci. Lett.* 147, 125-139.
- Monaco, C., Tortorici, L., Nicolich, R., Cernobori, L., Costa, M., 1996. From collisional to rifted basin: an example from the southern Calabrian arc (Italy). *Tectonophys.* 266, 233-249.
- Monaco, C., Tortorici, L., 2000. Active faulting in the Calabrian arc and eastern Sicily. *J. Geodyn.* 29, 407-424.
- Montenat, C., Barrier, P., d'Estevou, P., 1991. Some aspects of the recent tectonics in the Strait of Messina, Italy. *Tectonophys.* 194, 203-215.
- Montgomery, D.R., Brandon, M., T., 2002. Topographic controls on erosion rates in tectonically active mountain ranges, *Earth Pl. Sci. Lett.*, 201, 481-489.
- Montgomery, D.R., Dietrich, W.E., 1994. A physically-based model for the topographic control on shallow landsliding. *Water Resources Research* 30, 1153-1171.
- Morgagni, A., M. Marsigli, and E. Todini., 1993. Assessment of soil erosion in a number of small ephemeral streams in Calabria and Sicily. In Proc. Workshop on Soil Erosion in Semi-arid Mediterranean Areas, 137-148. European Society for Soil Conservation, Centro Studi per l'Economia applicata all'Ingegneria.
- Nandi, A., Shakoor, A., 2009. A GIS-based landslide susceptibility evaluation using bivariate and multivariate statistical analyses, *Eng. Geol.* 110, 11-20.

- Negredo, A.M., Sabadini, R., Bianco, G., Fernandez, M., 1999. Three-dimensionalmodelling of crustal motions caused by subduction and continental convergence in the central Mediterranean. *Geophys. J. Int.* 136, 261-274.
- Neri, G., Oliva, G., Orecchio, B., Presti, D., 2006.A possible seismic gap within the highly seismogenic belt crossing Calabria and Eastern Sicily, Italy. *Bull. Seism. Soc. Am.* 96, 1321-1331.
- Ogniben, L., 1973, Schema geologico della Calabria in base ai dati odierni: *Geologica Romana* 12, 243-585.
- Ohlmacher, C.G., Davis, C.J., 2003. Using multiple regression and GIS technology to predict landslide hazard in northeast Kansas, USA. *Eng. Geol.* 69, 331-343.
- Oztekin, B., Topal, T., 2005. GIS-based detachment susceptibility analyses of a cut slope in limestone, Ankara—Turkey. *Environ. Geol.* 49, 124-132.
- Pachauri, A.K., Gupta, P.V., Chander, R., 1998. Landslide zoning in a part of the Garhwal Himalayas. *Environ. Geol.* 36 (3-4), 325-334.
- Palmstrom, A., 1982. The Volumetric Joint Count – A Useful and Simple Measure of the Degree of Rock Jointing, Proc 4th Int. Congress Int. Ass. Engng. Geol., New Delhi 5, 221-228.
- Panza, G.F., Raykova, R.B., Carminati, E., Doglioni, C., 2007. Upper mantle flow in the western Mediterranean. *Earth Planet. Sci. Lett.* 257, 200-214.
- Pellegrino, A., Prestininzi, A., 2007. Impact of weathering on the geomechanical properties of rocks along thermal-metamorphic contact belts and morpho-evolutionary processes: The deep-seated gravitational slope deformations of Mt. Granieri-Salincriti (Calabria– Italy), *Geomorphology* 87, 176-195.
- Piacentini D., Troiani, F., Soldati, M., Notarnicola, C., Savelli, D., Schneiderbauer, S., Strada, C., 2012. Statistical analysis for assessing shallow-landslide susceptibility in South Tyrol (south-eastern Alps, Italy), *Geomorphology* 151/152, 196-206.

Bibliografia

- Poli, S., Sterlacchini, S., 2007. Landslide Representation Strategies in Susceptibility Studies using Weights-of-Evidence Modeling Technique, *Nat. Resources Res.*, 16, 121-134.
- Polonia, A., Torelli, L., Gasperini, L., Mussoni, P., 2012. Active faults and historical earthquakes in the Messina Straits area (Ionian Sea). *Nat. Hazards Earth Syst. Sci.* 12, 2311-2328.
- Pradhan, B., Lee, S., 2010. Delineation of landslide hazard areas on Penang Island, Malaysia, by using frequency ratio, logistic regression, and artificial neural network models. *Environ. Earth Sci.* 60, 1037-1054.
- Priest, S.D.; Hudson, J.A., 1981. Estimation of discontinuity spacing and tracing length using scanline survey. *Int. J. Rock Mech. Min. Sci. Geomech. Ab.* 18, 183-197.
- Ray, R.L., De Smedt, F., 2009. Slope stability analysis on a regional scale using GIS: a case study from Dhading, Nepal. *Environ. Geol.* 57, 1603-1611.
- Rickenmann, D., 1999 Empirical Relationships for Debris Flows. *Nat. Hazards* 19, 47–77.
- Riebe, C.S., Kirchner, J.W., Granger, D.E., Finkel, R.C., 2001. Strong tectonic and weak climatic control of long-term chemical weathering rates, *Geology* 29(6), 511-514.
- Rodriguez, C.E., Bommer, J.J., Chandler, R.J., 1999. Earthquake-induced landslides: 1980-1997. *Soil Dynamics and Earthquake Engineering* 18, 325-346.
- Rossi, M., Guzzetti, F., Reichenbach, P., Mondini, A:C., Peruccacci, S., 2010. Optimal landslide susceptibility zonation based on multiple forecast. *Geomorphology* 114, 129-142.
- Ruff, M., Czurda, K., 2008. Landslide susceptibility analysis with a heuristic approach in the Eastern Alps (Vorarlberg, Austria), *Geomorphology* 94, 314-324.
- Ruff, M., Rohn, J., 2008. Susceptibility analysis for slides and rockfall: an example from the Northern Calcareous Alps (Vorarlberg, Austria), *Environ. Geol.* 55, 441-452.
- Rust, D., Kershaw, S., 2000. Holocene tectonic uplift patterns in northeastern Sicily: evidence from marine notches in coastal outcrops. *Marine Geol.* 167, 105-126.

Bibliografia

- Sabato, L., Tropeano, M., 2004. Fiumara: a kind of high hazard river, *Phys. Chem. Earth.* 29, 717-715.
- Salciarini, D., Godt, J.W., Savage, W.Z., Baum, R.L., Conversini, P., 2008. Modeling landslide recurrence in Seattle, Washington, USA, *Eng. Geol.* 102, 227-237.
- Salciarini, D., Godt, J.W., Savage, W.Z., Conversini, P., Baum, R.L., Michael, J.A., 2006. Modeling regional initiation of rainfall-induced shallow landslides in the eastern Umbria Region of central Italy, *Landslides* 3, 181-194.
- Santangelo, M., Gioia, D., Fiorucci, F., Cardinali, M., Guzzetti, F., Schiattarelli, M. 2011. Relationships between mass movements and tectonic uplift in the upper Sinni River valley, Calabria-Lucania border, Southern Italy, *Geophys. Res. Abstracts*, 13, 11674.
- Schellart, W.P., 2010. Mount Etna-Iblean volcanism caused by roll-back induced upper mantle upwelling around the Ionian slab edge: An alternative to the plume model. *Geol.* 38/8, 691-694
- Schellart, W.P., Freeman, J., Stegman, D.R., Moresi, L., May, D., 2007. Evolution and diversity of subduction zones controlled by slab width. *Letters to Nature* 446, 308-311.
- Schulz, W.H., Lidke, D.J., Godt, J.W.: 2008. Modeling the spatial distribution of landslide-prone colluvium and shallow groundwater on hillslopes of Seattle, WA. *Earth Surf. Process. Landforms* 33, 123-141.
- Scicchitano, G., Spampinato, C.R., Ferranti, L., Antonioli, F., Monaco, C., Capano, M., Lubritto, C., 2011. Uplifted Holocene shorelines at Capo Milazzo (NE Sicily, Italy): Evidence of co-seismic and steady-state deformation. *Q. Int.* 232, 201-213.
- Serpelloni, E., Anzidei, M., Baldi, P., Casula, g., Galvani, A., 2005. Crustal velocity and strain-rate fields in Italy and surrounding regions: new results from the analysis of permanent and non-permanent GPS networks. *Geophys. J. Int.* 161, 861-880.

- Serpelloni, E., Vannucci, G., Pondrelli, S., Argnani, A., Casula G., Anzidei, M., Baldi, P., Gasperini, P., 2007. Kinematics of the Western Africa-Eurasia plate boundary from focal mechanisms and GPS data. *Geophys. J. Int.* 169, 1180-1200.
- Serpelloni, E., Bürgmann, R., Anzidei, M., Baldi, P., Mastrolemo Ventura, B., Boschi, E., 2010. Strain accumulation across the Messina Straits and kinematics of Sicily and Calabria from GPS data and dislocation modeling. *Earth Pl. Sci. Lett.* 298, 347-360.
- Shorey, P.R., Barat, D., Das, M.N., Mukherjee, K.P., Singh, B., 1984. Schmidt hammer rebound data for estimation of large scale in situ coal strength. *Int. J. Rock Mech. Min. Sci. Geomech. Abstr.* 21, 39-42.
- Singh, T., Awasthi, A.K., 2010. Stream profiles as indicator of active tectonic deformation along the Intra-Foreland Thrust, Nahan Salient, NW India. *Current Sci.*, 98(1-10), 95-98.
- Singh, L.P. van Westen, C.J., Champati Ray, P.K., Pasquall, P, 2005. Accuracy assessment of InSAR derived inputmaps for landslide susceptibility analysis: a case study from the Swiss Alps, *Landslides* 2, 221-228.
- Snyder, N.P., Whipple, K. X., Tucker, G. E., Merritts, D. J., 2002. Interaction between onshore bedrock-channel incision and nearshore wave-base erosion forced by eustasy and tectonics, *Basin Res.* 14, 105-127.
- Snyder, N.P., Whipple, K. X., Tucker, G. E., Merritts, D. J., 2003. Channel response to tectonic forcing: field analysis of stream morphology and hydrology in the Mendocino triple junction region, northern California, *Geomorphology* 53, 97-127.
- Somma, R., Messina, A., Mazzoli, S., 2005. Syn-orogenic extension in the Peloritani Alpine Thrust Belt (NE Sicily, Italy): Evidence from the Ali Unit. *C. R. Geosci.* 337, 861-871.
- Spampinato, C.R., Scicchitano, G., Ferranti, L., Monaco, C. 2012. Raised Holocene paleo-shorelines along the Capo Schisò coast, Taormina: New evidence of recent co-seismic deformation in northeastern Sicily (Italy). *J. Geodyn.* 55, 18-31.

Bibliografia

- Stewart, I.S., Cundy, A., Kershaw, S., Firth, C., 1997. Holocene coastal uplift in the Taormina area, northeastern Sicily: implications for the southern prolongation of the calabrian seismogenic belt. *J. Geodyn.* 24(14), 37-50.
- Strahler, A.N., 1952. Hypsometric (area-altitude) analysis of erosional topography. *Geol. Soc. Am. Bull.* 63, 1117-1142.
- Strahler, A.N., 1958. Dimensional analysis applied to fluvially eroded landforms. *Geol. Soc. Am. Bull.* 69, 279-300.
- Süzen, M.L., Doyuran, V., 2004. A comparison of the GIS based landslide susceptibility assessment methods: multivariate versus bivariate, *Environ. Geol.* 45, 665-679.
- Troiani, F, Della Setta, M., 2008. The use of the Stream Length–Gradient index in morphotectonic analysis of small catchments: A case study from Central Italy. *Geomorphology* 102, 159-168.
- Tortorici, L., Monaco, C., Tansi, C., Cocina, O., 1995. Recent and active tectonics in the Calabrian arc (Southern Italy). *Tectonophys.* 243, 37-55.
- USGS (U.S. Geological Survey), 2004. Landslide Types and Processes. Fact Sheet 2004-3072.
- van Westen, C.J., 1997. Statistical landslide hazard analysis. In: Application guide, ILWIS 2.1 for Windows. ITC, Enschede, The Netherlands, 73-84.
- van Westen, C.J., Castellanos, E., Kuriakose, S.L., 2008. Spatial data for landslide susceptibility, hazard, and vulnerability assessment: An overview, *Eng. Geol.* 102, 112-131.
- van Westen, C.J., Rengers; N., Soeters, R., 2003. Use of geomorphological information in indirect landslide susceptibility assessment, *Nat. Haz.* 30, 399-419.
- van Westen, C.J., van Asch, T.W.J., Soeters, R., 2006. Landslide hazard and risk zonation—why is it still so difficult?, *Bull. Eng. Geol. Environ.* 65, 167-184.
- van Westen, C.J., Rengers, N., Terlien, M.T.J., Soeters, R., 1997. Prediction of the occurrence of slope instability phenomena through GIS-based hazard zonation, *Geol. Rundsch.* 86, 404-414.

- Varnes, D.J., 1978. Slope movements, types and processes. In: Schuster, R.L., Krizek, R.J. (eds), *Landslide analysis and control*, National Academy Sciences, Washington DC, 11-33.
- Vecchio, A.: Analisi delle caratteristiche idrauliche del bacino idrografico del T. Giampilieri: valutazione del “rialzo idraulico” in sezioni ostruite (evento 1 Ottobre 2009). University of Catania, 2011.
- Waltham, T., 2002. Unloading joints and rockfalls in Norway’s fiordlands. *Geology Today*, 18(6), 220-225.
- Wang, W.D., Xie, C.M., Du, X.G., 2009. Landslides susceptibility mapping based on geographical information system, GuiZhou, south-west China. *Environ. Geol.* 58, 33-43.
- Westaway, R., 1993. Quaternary uplift of Southern Italy. *J. Geophys. Res.* 98, 21741-21772.
- Wortel, M.J.R., Spakman, W., 2000. Subduction and slab detachment in the Mediterranean-Carpathian region. *Science* 290, 1910-1917.
- Wu, C.H., Chen, S.C., 2009. Determining landslide susceptibility in Central Taiwan from rainfall and six site factors using the analytical hierarchy process method. *Geomorphology* 112, 190-204.
- Xu, P., Grasso, A., Mahtab, 1990. Use of Schmidthammer for estimating mechanical properties of weak rock 6th Int. IAEG Congress, Balkema, Rotterdam, 511-519.
- Yalcin, A., 2008. GIS-based landslide susceptibility mapping using analytical hierarchy process and bivariate statistics in Ardesen (Turkey): Comparisons of results and confirmations. *Catena* 72, 1-121.
- Yale, D.P., 2003. Fault and stress magnitude controls on variations in the orientation of in situ stress, *Geol. Soc. Lond. Spec. Publ.*, 209, 55-64.